

# ESTIMATE OF ISODIAMETRIC CONSTANT FOR CLOSED SURFACES

TAKASHI SHIOYA

ABSTRACT. We give an explicit estimate of the area of a closed surface by the diameter and a lower bound of curvature. This is better than Calabi-Cao's estimate [2] for a nonnegatively curved two-sphere.

## 1. INTRODUCTION

We are interested in the relation between the area  $V$  and the diameter  $D$  of a closed two-dimensional Riemannian manifold  $M$  with a lower bound of curvature  $\kappa$ . It is a famous conjecture due to A. D. Alexandrov [1] that, if  $\kappa = 0$ , then

$$\frac{V}{D^2} \leq \frac{\pi}{2}$$

and the equality holds only if  $M$  is the double of a flat disk. This conjecture seems to be very hard to solve. Since Alexandrov raised the conjecture in 1955, there were several attempts to approach it (see [2, 5, 12, 13, 15, 16]), but it is still open. A trivial upper bound of  $V/D^2$  is  $\pi$ , which follows from the Bishop volume comparison theorem. For a (not necessarily zero) lower bound  $\kappa$  of curvature, we have  $V \leq v_\kappa(D)$ , where  $v_\kappa(r)$  denotes the area of an  $r$ -ball in a complete simply connected two-dimensional space form with constant curvature  $\kappa$ , i.e.,

$$v_\kappa(r) = \begin{cases} \pi r^2 & \text{if } \kappa = 0, \\ 2\pi\kappa^{-1}(1 - \cos \sqrt{\kappa}r) & \text{if } \kappa > 0, \\ 2\pi\kappa^{-1}(1 - \cosh \sqrt{-\kappa}r) & \text{if } \kappa < 0. \end{cases}$$

Let  $R$  be the *radius* of  $M$ , i.e.,

$$R := \inf_{p \in M} \sup_{q \in M} d(p, q).$$

Note that  $D/2 \leq R \leq D$ . The Bishop volume comparison theorem also implies

$$(1.1) \quad V \leq v_\kappa(R).$$

---

*Date:* August 1, 2014.

*2010 Mathematics Subject Classification.* Primary 53C20.

*Key words and phrases.* A. D. Alexandrov's conjecture, isodiametric inequality, area, diameter, curvature.

The author is partially supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science.

This gives a good estimate of  $V/D^2$  only in the case where  $R$  is far less than  $D$ . In this paper, we give a new estimate of  $V/D^2$  which is effective for  $R$  close to  $D$  (see Proposition 3). Combining this with (1.1), we obtain the following main theorem.

**Theorem 1.** *Let  $M$  be a two-dimensional closed Riemannian manifold with area  $V$ , diameter  $D$ , Euler number  $\chi$ , and with a lower Gaussian curvature bound  $\kappa$ , and let  $k := \kappa D^2$ . We define*

$$w(k) := \frac{v_k(1)}{\pi} = \begin{cases} 1 & \text{if } k = 0, \\ 2k^{-1}(1 - \cos \sqrt{k}) & \text{if } k > 0, \\ 2k^{-1}(1 - \cosh \sqrt{-k}) & \text{if } k < 0, \end{cases}$$

$$\lambda_\chi(k) := 2\chi - k w(k),$$

$$\alpha_\chi(k) := \begin{cases} w(k) + 360^{-1}k \lambda_\chi(k) & \text{if } k \geq 0, \\ w(k)^2 & \text{if } k < 0. \end{cases}$$

Then we have

$$\frac{V}{D^2} \leq v_k \left( \sqrt{\frac{-6w(k) + 6\sqrt{w(k)^2 + 3^{-1}\lambda_\chi(k)\alpha_\chi(k)}}{\lambda_\chi(k)}} \right),$$

provided  $\lambda_\chi(k) > 0$ .

$\lambda_\chi(k)$  is a quantity of how much the surface is far from the space form. Note in fact that we always have  $\lambda_\chi(k) \geq 0$  and that  $\lambda_\chi(k) = 0$  holds if and only if  $M$  has *constant* nonnegative Gaussian curvature (see Proposition 5), for which we know the exact range of  $V/D^2$  (see Remark 6). Taking the limit of the estimate in the theorem as  $\lambda_\chi(k) \rightarrow 0+$  implies that  $V/D^2 \leq v_k(1)$  for  $\lambda_\chi(k) = 0$ , which is optimal in the case of constant positive curvature.

The right-hand side of the inequality in the theorem is continuous in  $k$  even at  $k = 0$ .

Letting  $\kappa := 0$  in the theorem yields the following.

**Corollary 2.** *Let  $M$  be a two-dimensional Riemannian manifold with nonnegative Gaussian curvature such that  $M$  is homeomorphic to either a two-sphere  $S^2$  or a real projective plane  $\mathbb{R}P^2$ . Then we have*

$$\frac{V}{D^2} \leq \frac{\sqrt{9+6\chi}-3}{\chi}\pi < \begin{cases} 2.486 & \text{if } M \simeq S^2, \\ 2.743 & \text{if } M \simeq \mathbb{R}P^2. \end{cases}$$

In the nonnegative curvature case, there were a few results for the estimate of  $V/D^2$  [2, 13, 15]. For  $M \simeq S^2$ , the best one that I had ever known was  $8/\pi \doteq 2.548$ , which is due to Calabi and Cao [2], where their estimate follows from the first nonzero eigenvalue estimate by Hirsch [11] and Zhong-Yang [17]. Our estimate, Corollary 2, is better than Calabi-Cao's estimate. For  $M \simeq \mathbb{R}P^2$ , Corollary 2 is also better than

the estimate in our previous paper [15]. The idea of the proof in this paper is different from any former results.

## 2. ESTIMATE BY RADIUS

The purpose of this section is to prove the following proposition, which is the key to the proof of the main theorem.

**Proposition 3.** *With the notations as in Theorem 1, we have*

$$V \leq \begin{cases} \pi D^2 - \frac{2\pi \chi \tilde{v}_\kappa(R)}{v_\kappa(D)} & \text{if } \kappa \geq 0, \\ \frac{\pi D^2 - 2\pi \chi \tilde{v}_\kappa(R)/v_\kappa(D) - \kappa \tilde{v}_\kappa(D)}{1 - \kappa \tilde{v}_\kappa(R)/v_\kappa(D)} & \text{if } \kappa < 0, \end{cases}$$

where  $R$  denotes the radius of  $M$  and

$$\begin{aligned} \tilde{v}_\kappa(r) &:= \int_0^r \int_0^t v_\kappa(s) ds dt \\ &= \begin{cases} \pi r^4/12 & \text{if } \kappa = 0, \\ \pi \kappa^{-2}(\kappa r^2 + 2 \cos(\sqrt{\kappa} r) - 2) & \text{if } \kappa > 0, \\ \pi \kappa^{-2}(\kappa r^2 + 2 \cosh(\sqrt{-\kappa} r) - 2) & \text{if } \kappa < 0. \end{cases} \end{aligned}$$

As a direct consequence of Proposition 3, we have the following.

**Corollary 4.** *Let  $M$  be a two-dimensional closed Riemannian manifold with nonnegative Gaussian curvature. Then we have*

$$\frac{V}{D^2} \leq \pi \left(1 - \frac{\chi}{6} \rho^4\right),$$

where  $\rho := R/D$ .

*Proof of Proposition 3.* We set, for  $p \in M$ ,

$$R_p := \sup_{q \in M} d(p, q),$$

where  $d$  denotes the Riemannian distance function on  $M$ . Note that  $R_p$  is continuous in  $p \in M$ . Let  $B(p, r)$  indicate the  $r$ -ball centered at  $p \in M$  and  $\partial B(p, r)$  its boundary. Denote by  $\kappa(p, r)$  the sum of the total geodesic curvature of  $\partial B(p, r)$  and of the exterior angles of  $B(p, r)$  at cut points in  $\partial B(p, r)$  from  $p$ , which is well-defined a.e.  $r \in (0, R_p)$ . By Fiala-Hartman's discussion (see [4, 10] and [14, Theorem 4.4.1]), the length of  $\partial B(p, r)$ , say  $L(p, r)$ , satisfies

$$\frac{d}{dr} L(p, r) \leq \kappa(p, r)$$

for a.e.  $r \in (0, R_p)$ . It follows from the Gauss-Bonnet theorem that

$$\kappa(p, r) \leq 2\pi - \int_{B(p, r)} K(q) dq$$

for a.e.  $r \in (0, R_p)$ , because the Euler number of a proper subset of  $M$  is at most one, where  $K$  denotes the Gaussian curvature function on  $M$  and  $dq$  the area (volume) measure with respect to the variable  $q \in M$ . Note that  $\int_{B(p,r)} K(q) dq$  is continuous in  $(p, r) \in M \times [0, +\infty)$ . We therefore have

$$V = \int_0^{R_p} L(p, t) dt \leq \iint_{0 \leq s \leq t \leq R_p} \left( 2\pi - \int_{B(p,s)} K(q) dq \right) ds dt,$$

where we note that the right-hand side is continuous in  $p \in M$ . Integrating the both sides of the above inequality with respect to  $p \in M$ , we have, by  $R_p \leq D$ ,

$$(2.1) \quad V^2 \leq \iiint_{\substack{0 \leq s \leq t \leq D \\ p \in M}} 2\pi ds dt dp - I = \pi D^2 V - I,$$

where

$$I := \iiint_{\substack{0 \leq s \leq t \leq R_p \\ d(p,q) < s}} K(q) ds dt dp dq.$$

We are going to estimate  $I$  from below.

We assume  $\kappa \geq 0$ . Denote by  $V(q, s)$  the area of the ball  $B(q, s)$ . Since  $K \geq 0$ ,  $R_p \geq R$ , and by the Bishop-Gromov volume comparison theorem, we see that

$$\begin{aligned} I &\geq \iiint_{\substack{0 \leq s \leq t \leq R \\ d(p,q) < s}} K(q) ds dt dp dq = \iint_{\substack{0 \leq s \leq t \leq R \\ q \in M}} K(q) V(q, s) ds dt dq \\ &\geq \frac{V}{v_\kappa(D)} \iint_{\substack{0 \leq s \leq t \leq R \\ q \in M}} K(q) v_\kappa(s) ds dt dq \end{aligned}$$

and by the Gauss-Bonnet theorem,

$$= \frac{2\pi\chi V}{v_\kappa(D)} \iint_{0 \leq s \leq t \leq R} v_\kappa(s) ds dt = \frac{2\pi\chi \tilde{v}_\kappa(R)}{v_\kappa(D)} V.$$

Combining this with (2.1) yields

$$V^2 \leq \pi D^2 V - \frac{2\pi\chi \tilde{v}_\kappa(R)}{v_\kappa(D)} V,$$

which implies

$$V \leq \pi D^2 - \frac{2\pi\chi \tilde{v}_\kappa(R)}{v_\kappa(D)}.$$

We next assume  $\kappa < 0$ . Since  $K - \kappa \geq 0$  and  $R \leq R_p \leq D$ ,

$$\begin{aligned} I &\geq \iiint\limits_{\substack{0 \leq s \leq t \leq R \\ d(p,q) < s}} (K(q) - \kappa) ds dt dp dq + \iiint\limits_{\substack{0 \leq s \leq t \leq D \\ d(p,q) < s}} \kappa ds dt dp dq \\ &= \iiint\limits_{\substack{0 \leq s \leq t \leq R \\ q \in M}} (K(q) - \kappa) V(q, s) ds dt dq + \iiint\limits_{\substack{0 \leq s \leq t \leq D \\ q \in M}} \kappa V(q, s) ds dt dq \end{aligned}$$

and by the Bishop-Gromov and Bishop inequalities,

$$\begin{aligned} &\geq \frac{V}{v_\kappa(D)} \iiint\limits_{\substack{0 \leq s \leq t \leq R \\ q \in M}} (K(q) - \kappa) v_\kappa(s) ds dt dq + \iiint\limits_{\substack{0 \leq s \leq t \leq D \\ q \in M}} \kappa v_\kappa(s) ds dt dq \\ &= \frac{\tilde{v}_\kappa(R)}{v_\kappa(D)} (2\pi\chi - \kappa V) V + \kappa \tilde{v}_\kappa(D) V, \end{aligned}$$

which together with (2.1) leads us to the proposition. This completes the proof.  $\square$

### 3. ESTIMATE BY DIAMETER

**Proposition 5.** *With the notations as in Theorem 1, we have*

$$\lambda_\chi(k) \geq 0$$

*and the equality holds if and only if  $M$  has constant nonnegative Gaussian curvature.*

*Proof.* We first assume  $k > 0$ . Then,  $\chi = 2$  (i.e.,  $M \simeq S^2$ ) or  $\chi = 1$  (i.e.,  $M \simeq \mathbb{R}P^2$ ). Note that  $\lambda_\chi(k) = 2(\chi - 1 + \cos \sqrt{k})$ .

In the case where  $\chi = 2$ , we have  $\lambda_\chi(k) \geq 0$ , and the equality holds if and only if  $k = \pi^2$ , which is equivalent to  $M$  being a two-sphere with positive constant curvature by the Toponogov maximal diameter theorem [3, Theorem 6.5].

In the case where  $\chi = 1$ , we have  $\lambda_\chi(k) \geq 0$  (resp.  $= 0$ ) if and only if  $k \leq \pi^2/4$  (resp.  $= \pi^2/4$ ). The theorem in this case follows from the Grove-Shiohama diameter sphere theorem [9] and the Gromoll-Grove diameter rigidity theorem [6].

We next assume  $k \leq 0$ . By the Gauss-Bonnet theorem and the Bishop volume comparison theorem,

$$\pi \lambda_\chi(k) = \int_M K(p) dp - \kappa v_\kappa(D) \geq \kappa (V - v_\kappa(D)) \geq 0.$$

If  $\lambda_\chi(k) = 0$  and  $k < 0$ , then  $K \equiv \kappa$  and  $V = v_\kappa(D)$ , so that the cut-locus at any point  $p$  in  $M$  is contained in  $\partial B(p, D)$ , which is a

contradiction (cf. [7]). Therefore, if  $\lambda_\chi(k) = 0$ , then  $k = 0$ , which implies

$$\int_M K(p) dp = 2\pi\chi = \pi\lambda_\chi(k) = 0$$

and so  $M$  has a flat metric. This completes the proof.  $\square$

*Remark 6.* If a closed two-manifold has constant nonnegative Gaussian curvature, then it is isometric to either a two-sphere with a metric of constant positive curvature, a real projective plane with a metric of constant positive curvature, a two-torus with a flat metric, or a Klein bottle with a flat metric. It is easy to see that  $V/D^2 \leq 2$  for a closed two-manifold with a flat metric. For a two-sphere with a metric of constant curvature,  $V/D^2 = 4/\pi$ . For a real projective plane with a metric of constant curvature,  $V/D^2 = 8/\pi$ .

*Proof of Theorem 1.* Assume  $\lambda_\chi(k) > 0$ . Scaling the metric of  $M$ , we may assume that  $D = 1$ . Then,  $k = \kappa$  and  $1/2 \leq R \leq 1$ .

We first prove the theorem in the case where  $k \geq 0$ . Note that  $\chi \geq 0$  in this case. Combining Proposition 3 with (1.1) implies

$$\frac{V}{D^2} \leq \sup_{1/2 \leq r \leq 1} \min\{ \pi - 2\pi\chi \tilde{v}_k(r)/v_k(1), v_k(r) \}.$$

By considering the monotonicity of  $\pi - 2\pi\chi \tilde{v}_k(r)/v_k(1)$  and  $v_k(r)$  in  $r$ , the supremum above is attained at a unique solution to the equation

$$(3.1) \quad \pi - 2\pi\chi \tilde{v}_k(r)/v_k(1) = v_k(r).$$

We will estimate the solution  $r$  from above. For simplicity we set  $f(r) := \tilde{v}_k(r)/\pi$ . Since  $w(k) = v_k(1)/\pi$  and  $v_k(r)/\pi = r^2 - k f(r)$ , the equation (3.1) implies

$$\lambda_\chi(k) f(r) = w(k) - w(k) r^2.$$

It follows that

$$f(r) \geq \frac{r^4}{12} - \frac{k r^6}{360} \geq \frac{r^4}{12} - \frac{k}{360},$$

so that

$$\lambda_\chi(k) \left( \frac{r^4}{12} - \frac{k}{360} \right) \leq w(k) - w(k) r^2,$$

which implies

$$(3.2) \quad \frac{1}{12} \lambda_\chi(k) r^4 + w(k) r^2 - \alpha_\chi(k) \leq 0.$$

We therefore have

$$r^2 \leq \frac{-6 w(k) + 6 \sqrt{w(k)^2 + 3^{-1} \lambda_\chi(k) \alpha_\chi(k)}}{\lambda_\chi(k)},$$

which together with  $V \leq v_k(r)$  proves the theorem in this case.

We next prove the theorem in the case where  $k < 0$ . Since the derivative

$$\frac{d}{d\tilde{v}} \left( \frac{\pi - 2\pi\chi \tilde{v}/v_k(1) - k \tilde{v}_k(1)}{1 - k \tilde{v}/v_k(1)} \right) = - \frac{\lambda_\chi(k)}{(1 - k \tilde{v}/v_k(1))^2 w(k)}.$$

is negative, the function

$$\frac{\pi - 2\pi\chi \tilde{v}_k(r)/v_k(1) - k \tilde{v}_k(1)}{1 - k \tilde{v}_k(r)/v_k(1)}$$

is monotone decreasing in  $r$ . In the same way as before, we have  $V \leq v_k(r)$  for a unique solution  $r$  to the equation

$$(3.3) \quad \frac{\pi - 2\pi\chi \tilde{v}_k(r)/v_k(1) - k \tilde{v}_k(1)}{1 - k \tilde{v}_k(r)/v_k(1)} = v_k(r).$$

For the solution  $r$ , we have

$$\pi - 2\pi\chi \tilde{v}_k(r)/v_k(1) - k \tilde{v}_k(1) \geq v_k(r)$$

and then

$$\lambda_\chi(k)f(r) + w(k)r^2 - w(k)^2 \leq 0,$$

which together with  $f(r) \geq r^4/12$  yields (3.2). This completes the proof of Theorem 1.  $\square$

*Remark 7.* In the case where  $k < 0$ , we have, by (3.3),

$$k^2 f(r)^2 + (\lambda_\chi(k) - k r^2) f(r) + w(k) r^2 - w(k)^2 = 0,$$

which together with  $f(r) \geq r^4/12$  implies a quartic inequality for  $r^2$ , leading to a better and much more complex form of the estimate of  $V/D^2$ .

## REFERENCES

- [1] A. D. Alexandrow, *Die innere Geometrie der konvexen Flächen*, Akademie-Verlag, Berlin, 1955 (German).
- [2] E. Calabi and J. G. Cao, *Simple closed geodesics on convex surfaces*, J. Differential Geom. **36** (1992), no. 3, 517–549.
- [3] J. Cheeger and D. G. Ebin, *Comparison theorems in Riemannian geometry*, AMS Chelsea Publishing, Providence, RI, 2008. Revised reprint of the 1975 original.
- [4] F. Fiala, *Le problème des isopérimètres sur les surfaces ouvertes à courbure positive*, Comment. Math. Helv. **13** (1941), 293–346 (French).
- [5] P. Freitas and D. Krejčířík, *Alexandrov's isodiametric conjecture and the cut locus of a surface*. preprint, arXiv:1406.0811v1.
- [6] D. Gromoll and K. Grove, *A generalization of Berger's rigidity theorem for positively curved manifolds*, Ann. Sci. École Norm. Sup. (4) **20** (1987), no. 2, 227–239.
- [7] K. Grove and P. Petersen, *Manifolds near the boundary of existence*, J. Differential Geom. **33** (1991), no. 2, 379–394.
- [8] K. Grove and P. Petersen V, *Volume comparison à la Aleksandrov*, Acta Math. **169** (1992), no. 1-2, 131–151, DOI 10.1007/BF02392759.
- [9] K. Grove and K. Shiohama, *A generalized sphere theorem*, Ann. of Math. (2) **106** (1977), no. 2, 201–211.

- [10] P. Hartman, *Geodesic parallel coordinates in the large*, Amer. J. Math. **86** (1964), 705–727.
- [11] J. Hersch, *Quatre propriétés isopérimétriques de membranes sphériques homogènes*, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A1645–A1648 (French).
- [12] J. Krysiak and Z. McCoy, *Alexandrov’s conjecture: On the intrinsic diameter and surface area of convex surfaces*. preprint.
- [13] T. Sakai, *On the isodiametric inequality for the 2-sphere*, Geometry of manifolds (Matsumoto, 1988), Perspect. Math., vol. 8, Academic Press, Boston, MA, 1989, pp. 303–315.
- [14] K. Shiohama, T. Shioya, and M. Tanaka, *The geometry of total curvature on complete open surfaces*, Cambridge Tracts in Mathematics, vol. 159, Cambridge University Press, Cambridge, 2003.
- [15] T. Shioya, *Diameter and area estimates for  $S^2$  and  $\mathbf{P}^2$  with non-negatively curved metrics*, Progress in differential geometry, Adv. Stud. Pure Math., vol. 22, Math. Soc. Japan, Tokyo, 1993, pp. 309–319.
- [16] V. A. Zalgaller, *A conjecture on convex polyhedra*, Sibirsk. Mat. Zh. **50** (2009), no. 5, 1070–1082, DOI 10.1007/s11202-009-0095-3 (Russian, with Russian summary); English transl., Sib. Math. J. **50** (2009), no. 5, 846–855.
- [17] J. Q. Zhong and H. C. Yang, *On the estimate of the first eigenvalue of a compact Riemannian manifold*, Sci. Sinica Ser. A **27** (1984), no. 12, 1265–1273.

MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, SENDAI 980-8578, JAPAN